Optimum Internal Medicine & Pediatrics
Dr. Christopher Burress
Board certified Internal Medicine and Pediatrics

Caring for your 9-month-old baby

Name_____ Weight_____ Height_____ Date_____

Keep Your Baby Safe

Use a Car Seat

Install the car seat properly in the back seat, rear-facing until your baby reaches the seat manufacturer's maximum rear-facing height or weight limit.

Use a Safe Crib

- Make sure the space between the slats is less than 2½ inches.
- Don't use a crib with sharp edges that your baby can hit.

Prevent Suffocation

- Put your baby on your baby's back to sleep.
- Don't let your baby play with balloons, rubber bands or objects with strings.
- Don't give your baby toys with small, removable parts (like buttons) or toys that fit completely in your baby's mouth. Give your baby toys made for your baby's age.

Prevent Injuries

- Don't leave your baby alone—ever. Use extra care when your baby is in water, near young children or pets, or on any surface from which your baby could fall.
- Don't use a walker that will let your baby move across the floor. Your baby does not need shoes until your baby can walk outdoors.
- Tell your health care provider if you have peeling paint in your home.
- "Child-proof" your home (see our "Child-Proofing Checklist").
- Set the water heater thermostat so that the water is not hotter than 120ºF.
- Make sure you have smoke detectors on your ceiling. Check twice a year to make sure they work.
- Keep your baby away from hot things (like cigarettes or an iron). Don't hold your baby when you are drinking something hot (like coffee) or near a hot stove or oven.
- Protect your baby from sunburn. Use sunscreen.
- **Never** shake, toss or hit your baby.

Provide Healthy Nutrition

- Breast milk or infant formula is all your baby needs. A healthy baby does not need vitamins. You may give solid foods. Start with iron-fortified rice cereal mixed with breast milk or formula.
- If you give other foods to your baby, give only one new food to your baby every few days. Single foods (like mashed carrots) are better than mixtures (like "chicken dinner"). Don't give your baby honey until your baby is at least 12 months old. Don't give your baby foods your baby might choke on (like nuts, beans, peas, grapes, and pieces of carrot or hot dog).
- Feed your baby at least 5 times daily. Avoid nighttime feedings.
- Don't prop a bottle in your baby's mouth. Don't put your baby in the crib with a bottle.

Optimum Internal Medicine & Pediatrics
Dr. Christopher Burress
Board certified Internal Medicine and Pediatrics

Practice Healthy Parenting and Care-giving

Encourage Healthy Development

- Create a schedule for your baby's day and try to keep to it.
- Excite your baby's sense of sight. Show your baby a sturdy, brightly colored picture book.
- Excite your baby's sense of touch. Let your baby touch objects with different textures (like soft, hard, glossy or furry).
- Excite your baby's sense of hearing. Talk and sing to your baby. Give your baby wooden spoons and plastic containers to play with.
- Strengthen your baby and your baby's coordination. Help your baby stack blocks and drink from a cup. Help your baby to crawl and to stand up.
- Encourage your baby socially. Play "pat-a-cake" and "peek-a-boo."
- **Never** yell at your baby. Keep your baby safe by moving your baby away from danger or re-directing your baby's attention to something else.
- Avoid screen time with your baby by limiting exposure to Cell Phones, Tablets, or Laptops.

Deal with Crying

- When your baby cries, ask yourself, "Is my baby hungry? Does your baby want to be changed? Does your baby want to be held? Is your baby too stimulated (like from too much touch, noise or light)? Is your baby in pain (like from teething)? Does your baby have a fever?"
- Treat the pain of teething with non-prescription teething medicine applied to the gum or with acetaminophen or ibuprofen.

Sleep

- Put your baby in the crib on your baby's back when your baby is drowsy but still awake.
- Your baby needs naps and a regular bedtime.
- Don't pick up your baby in the middle of the night. Comfort your baby with gentle strokes if your baby is crying.

Clean Your Baby

- If your baby boy is uncircumcised, don't try to pull back your baby's foreskin.
- Bathe your baby every other day. When you do, use warm water and mild baby soap.

Prevent Sickness—Identify It Early

- Do not induce vomiting or use ipecac. For suspected poisoning, call Poison Control at 1-800-222-1222. Save this number in your cell phone.
- Keep your baby away from cigarette smoke. Do not smoke with the baby in the car; do not smoke in the home.
- Call the clinic right away if your baby is coughing a lot or may be dehydrated. Sneezing, watery eyes and a stuffy or runny nose are cold symptoms. A fever is an under-the-arm temperature of 101°F or higher. A dehydrated baby has fewer than five wet diapers a day, is much less active than usual, or has dry lips and a pasty mouth.

Development

All babies develop at their own rate. At this age you may notice that your baby:

- Sits without support
- Tries to crawl, but might still be dragging legs or going backward
- Pulls on something to try to stand
- Takes steps while holding on to furniture
- Moves objects from hand to hand
- Picks up small objects with index finger and thumb
- Understands your baby's name, "no" and "bye-bye"
- Repeats syllables ("da-da" and "ba-ba")

Optimum Internal Medicine & Pediatrics
Dr. Christopher Burress
Board certified Internal Medicine and Pediatrics

- Waves, claps and copies others
- Enjoys social games such as peekaboo

Your baby's next visit is when your baby is at least 12-months-old. At this check-up your baby will receive more immunizations (shots). We may check for anemia or infection with the tuberculosis bacteria. Remember to bring your baby's shot record with you. Bring acetaminophen or ibuprofen for infants with you to the next visit.